

---

# Effects of manure injection tool type and tool spacing on soil nutrient levels and spring barley performance

B. Assefa<sup>1</sup>, Y. Chen<sup>1\*</sup>, K. Buckley<sup>2</sup> and W. Akinremi<sup>3</sup>

<sup>1</sup>Department of Biosystems Engineering, University of Manitoba, Winnipeg, Manitoba R3T 5V6, Canada; <sup>2</sup>Brandon Research Center, Agriculture & Agri-Food Canada, Brandon, Manitoba R7A 5Y3, Canada; and <sup>3</sup>Department of Soil Science, University of Manitoba, Winnipeg, Manitoba R3T 2N2, Canada. \*Email: ying\_chen@umanitoba.ca

---

Assefa, B., Chen, Y., Buckley, K. and Akinremi, W. 2006. **Effects of manure injection tool type and tool spacing on soil nutrient levels and spring barley performance.** Canadian Biosystems Engineering /Le génie des biosystèmes au Canada **48**: 2.45 - 2.54. A three-year field trial was conducted to study the effects of two manure injection tool types and three tool spacings on soil nutrient levels and crop response in a 2 x 3 factorial experiment. Liquid swine manure was injected using coulter and furrower injectors at 300- (S300), 600- (S600), and 900-mm (S900) tool spacings. Extractable soil NO<sub>3</sub>-N, NH<sub>4</sub>-N, P<sub>2</sub>O<sub>5</sub>, K, SO<sub>4</sub>-S, pH, and electrical conductivity (EC), plant number of tillers, heads, and main stem length, plant biomass, grain and straw yields, total N and P in plant biomass, grain, and straw were measured. Application of manure with a furrower proved to be advantageous over a coulter in many ways. Use of a furrower resulted in 40 to 60% higher soil NO<sub>3</sub>-N than a coulter at 0-300 mm soil depth at the time of rapid plant development in the second and third years of the experiment. Furthermore, use of a furrower resulted in 10% more plant biomass, 13, 3, and 16% higher total N in plant biomass, grain, and straw, respectively, and 2.5 and 13% higher total P in grain and straw, respectively, compared to use of a coulter in the first year of the experiment. Increased tool spacing decreased total N in plant biomass, grain, and straw. Soil nutrient levels also decreased with increase in tool spacing in one year of the study. In the other years, S300 resulted in higher soil NO<sub>3</sub>-N and NH<sub>4</sub>-N at 0-300 mm soil depth than S600. Plant number of tillers, heads, and main stem length in S300 were in some cases equivalent to and in other cases higher than those of S600 and S900. **Keywords:** manure, injection, tool, tool spacing, soil, nutrient, crop, yield.

Une étude échelonnée sur trois ans a été entreprise pour examiner l'influence de l'espacement entre les outils pour deux types d'enfouisseurs de lisier sur le placement de nutriments et la production des cultures, par le biais d'un plan expérimental factoriel 2 X 3. Le lisier de porc était enfouis directement dans le sol avec soit des enfouisseurs du type "coute à disque" ou bien du type "soc patte d'oie" (profil en "V") espacés de 300- (S300), 600- (S600) ou 900-mm (S900). Les niveaux labiles de NO<sub>3</sub>-N, NH<sub>4</sub>-N, P<sub>2</sub>O<sub>5</sub>, K, SO<sub>4</sub>-S, le pH et la conductivité électrique (EC), la population de tiges, le nombre de gerbes, ainsi que la longueur de la tige principale, la biomasse totale, les rendements de grain et paille, les niveaux N et P totaux dans la biomasse récoltée ont tous été mesurés. L'enfouissement du lisier avec un soc est plus efficace qu'avec l'enfouisseur à disque à plusieurs égards. L'enfouissement avec un soc a favorisé des niveaux 40 à 60% plus élevés de NO<sub>3</sub>-N que le coute à disque pour les premiers 300 mm du sol au moment coïncidant à une croissance rapide de la culture durant la deuxième et troisième année de l'étude. L'enfouisseur à soc a résulté en une biomasse 10% plus élevée, des niveaux de N totale

13%, 3% et 16% plus élevés dans la biomasse, le grain et la paille, respectivement, ainsi que un P total 2.5% et 13% plus important dans le grain et la paille durant la première année de l'étude. La quantité de N totale mesurée dans la biomasse, grain et paille, décroît avec l'espacement plus grand entre les enfouisseurs. Les niveaux de nutriments dans le sol ont aussi décrû avec l'écartement plus grand entre les enfouisseurs pour une année de l'étude. Les autres années, S300 a permis des niveaux plus élevés de NO<sub>3</sub>-N et NH<sub>4</sub>-N dans les premiers 300 mm du sol que S600. Le nombre de tiges et gerbes, ainsi que la longueur de la tige principale avec S300 étaient soit équivalentes ou plus élevées que pour S600 et S900. **Mots clés :** lisier, injection, enfouissement, outils, espacement, sol, nutriment, culture, rendement.

## INTRODUCTION

Land application of manure is considered the most economical management practice that enables recycling of the nutrients contained in manure. However, surface applications of liquid manure may lead to odour problems and nutrient losses due to surface runoff and ammonia volatilization. To maximize the returns from liquid manure and reduce environmental impact of applications, many livestock producers choose to inject the manure into the soil. Research findings also indicate that manure injection is preferable to surface application because it reduces odour, surface runoff, and loss of ammonia (Sutton 1994; Hoff et al. 1981), which eventually contributes to increased crop yields (Chen and Samson 2002; Mooleki et al. 2002).

Liquid manure injection involves selection of the right injection tool and tool spacing. Several types of injection tools, which include sweeps, discs, knives, chisels, and coulters, have been developed for injecting liquid manure below the soil surface. These tools are generally classified into two main groups: winged tools, such as furrowers and sweeps, and non-winged tools, such as discs, knives, and coulters. Winged tools place manure in wider bands and non-winged tools place manure in narrower bands (Rahman et al. 2004; Warner and Godwin 1988). Winged tools are more widely used compared to the non-winged ones because the former allow higher application rates and better soil-manure mixing (Chen and Tessier 2001).

Tool spacing determines the distance between manure bands. Narrow tool spacing may increase the capital cost of the



**Table 2. Dates of field operations and measurements.**

Field activity	Year		
	2002	2003	2004
Manure injection	May 28	June 17	June 22
Seeding	June 4	June 20	June 25
First soil sampling	June 18	July 8	July 14
Second soil sampling	July 2	July 29	August 4
Third soil sampling	July 16	August 19	November 4
Fourth soil sampling	July 30	NA*	NA
Fifth soil sampling	August 13	NA	NA
Plant sampling	August 13	August 19	September 8
Yield harvesting	September 5, 11	September 3, 4	October 16

\*NA = not applicable

### Field operation procedure

Injection was performed after tillage with a heavy duty field cultivator and prior to seeding in the spring of each growing season. A custom built distributor delivered manure through flexible hoses to the injection tools. Manure was injected using seven, four, and three tools mounted on the toolbar for the S300, S600, and S900 tool spacing treatments, respectively. Manure was injected to a depth of 100 mm at an average rate of 34 m<sup>3</sup>/ha for all plots and years. Manure flow rate from the tank was kept constant by maintaining a constant pumping rate during the entire manure injection operation. The travel speed of the injector was also kept constant. In all three years, plots were seeded to hulless spring barley (Cultivar: AC Bacon) a few days after manure injection. Dates for the field operations are given in Table 2.

### Measurements

**Soil and manure background** Immediately prior to manure injection, soil samples were collected from five random plots across the entire field sites to determine the soil moisture content and bulk density. The sampling was done to a depth of 150 mm using 52-mm diameter core samplers. Soil moisture content and bulk density were determined by oven drying for 24 h at 105°C. Manure samples were taken for analysis two weeks in advance of manure application. Electrical conductivity and pH measurements were performed on a 10:1 dilution of liquid manure with distilled water. Ammonia concentration in the diluted mixture was determined by an ion specific electrode against a certified standard. Moisture content was measured after oven drying to a constant weight at 105°C. Total nitrogen was measured by standard Kjeldahl analysis (AOAC 1990). Total P, K, Ca, Na, Mg, and S were measured by total digestion of the sample in nitric/perchloric acid and analysed by inductively-coupled plasma spectrometry.

**Soil nutrients** Soil samples for nutrient analysis were collected along transects perpendicular to the travel direction of the injector. In each plot, three transects were identified in three random locations. Along the transects, samples were collected from positions located at 0, 150, 300, and 450-mm distances from the centre line of a manure band. Soil samples were

collected five times to a depth of 300 mm in the growing season of 2002 and three times each growing season in 2003 and 2004 in two depth ranges (0-300 and 300-600 mm). The samples collected from the three locations per plot were mixed together depth-wise to form a composite sample of the respective position.

The soil samples were air dried and ground to less than 2-mm size prior to analyses. Samples collected in 2002 were analysed for extractable NO<sub>3</sub>-N, NH<sub>4</sub>-N, P<sub>2</sub>O<sub>5</sub>, K, SO<sub>4</sub>-S, and pH and electrical conductivity (EC). In 2003 the soil samples were analysed for extractable NO<sub>3</sub>-N, P<sub>2</sub>O<sub>5</sub>, and K. Soil samples collected in 2004 were analysed for extractable NO<sub>3</sub>-N, NH<sub>4</sub>-N, P<sub>2</sub>O<sub>5</sub>, and pH and EC. Methods used for analyses are presented in Table 3.

### Plant tillers, heads, main stem length, and above ground biomass

At the plants' soft dough stage, 20-40 plants were uprooted. The number of tillers and heads per plant were counted, and the main stem length of each plant was measured with a ruler. At the same time, a 500-mm wide plant strip was cut across each plot width to measure the amount of above ground biomass. The biomass samples were oven-dried at 60°C for 72 h (ASAE 2004a) to determine the dry matter of the plant biomass. Total N and P in the biomass were determined on digested samples of ground plant biomass by the standard acid (H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub>) digestion method described in Thomas et al. (1967). A Technicon Autoanalyzer (Technicon Corporation, Terrytown, NY) was used to calorimetrically determine total N and P in the digest samples.

**Grain and straw yield** A plot combine was used to harvest the plots for yield measurements in 2002 and 2003. Entire plots were harvested and the harvest areas were calculated after adjustment for crop removal for the biomass measurement. In 2004, harvesting was done by hand, as wet soil and lodged crop did not allow for using the plot combine. Crop samples were collected by cutting a 1-m<sup>2</sup> area at three random locations from each plot. Samples were threshed in the lab. Grain and straw samples were separately weighed, oven dried at 60°C for 72 h and weighed again to determine the dry matter of the grain and straw yields at 11% moisture content. Total N and P in the grain and straw were also determined the same way as for the ground biomass.

### Statistical analyses

The data were analysed separately under each year using SAS software (SAS Institute Inc. 2001). Analysis of variance was carried out using the general linear model procedure to determine the means of each variable. Standard errors were used to determine differences among treatment means. All comparisons were made at a probability of 0.1 because of the inherent high variability of soil. The analyses results revealed that there were no interactions between the experimental factors (tool type and tool spacing). The main effects of the factors on soil nutrient levels and crop response are presented in the following sections.

**Table 3. Methods used for soil analysis.**

Soil property	Analytical method and reference		
	2002	2003	2004
NO <sub>3</sub> -N	Cadmium reduction procedure (Maynard and Kalra 1993)	Sodium bicarbonate method (Olsen and Sommers 1982; Hamm et al. 1970)	Automated cadmium reduction method (Greenberg et al. 1992)
NH <sub>4</sub> -N	Automated phenate method (Greenberg et al. 1992)	Not analysed	Automated phenate method (Greenberg et al. 1992)
P	Modified Kelowna soil test (Asworth and Mrazek 1995)	Sodium bicarbonate method (Olsen and Sommers 1982; Hamm et al. 1970)	Modified Kelowna soil test (Qian et al. 1994)
K	Automated flame photometry (Alberta Research Council 1996)	Sodium bicarbonate method (Olsen and Sommers 1982; Hamm et al. 1970)	Not analysed
SO <sub>4</sub> -S	Automated methylthymol blue method (Clesceri et al. 1998)	Not analysed	Not analysed
pH	1:2 soil water extract (Hendershot et al. 1993)	Not determined	1:2 soil water extract (Hendershot et al. 1993)
EC	1:2 soil water extract (McKeague 1976)	Not determined	1:2 soil water extract (McKeague 1976)

**Table 4. Growing season monthly air temperature and precipitation for the three years and averages of 16 years prior to 2002.**

Growing season	Air temperature (°C)						Precipitation (mm)					
	April	May	June	July	August	Average	April	May	June	July	August	Total
2002	2	8	18	20	17	13	16	8	75	51	101	251
2003	5	12	16	20	22	15	45	42	65	5	28	185
2004	4	7	14	18	14	11	21	160	39	76	74	369
Average	4	12	17	19	18	14	24	60	71	77	58	291

**Table 5. Soil (150 mm deep) bulk density and moisture content at the time of manure injection and composition of the manure applied (wet basis).**

Term	Year		
	2002	2003	2004
<i>Soil property</i>			
Bulk density (Mg/m <sup>3</sup> )	0.80	0.85	0.80
Gravimetric moisture content(%)	24	34	36
<i>Manure characteristics</i>			
Total N (kg/ML)	2.40	2.90	3.50
Organic N (kg/ML)	0.20	0.60	0.40
NH <sub>4</sub> -N (kg/ML)	2.20	2.30	3.00
Nitrate and nitrite N (kg/ML)	0.10	0.10	0.10
Total P (kg/ML)	0.04	0.79	0.55
Total K (kg/ML)	1.32	1.51	1.84
Total S (kg/ML)	0.11	0.20	0.23
Solid content (%)	1.10	2.10	1.60
EC (dS/m)	15.85	18.10	19.80
pH	7.60	7.90	7.40

## RESULTS and DISCUSSION

### Weather conditions and soil and manure background

The monthly air temperature and precipitation data for the three growing seasons and averages of 16 years prior to 2002 are given in Table 4. The data were obtained from Brandon Research Centre weather station located within 1 km of the sites. The growing seasons of 2002 and 2003 were relatively dryer than the 16 year average whereas that of 2004 was the wettest and coldest of all.

Soil (150 mm deep) bulk density and moisture content and composition of the manure applied are presented in Table 5. At the time of manure application, the soils had low bulk densities as the measurement was done a few days after spring tillage. When averaged over three years, total N (2.9 kg /ML) and total P (0.6 kg /ML) in the manure were similar to and lower than the mean total N and mean total P in swine manure in Manitoba (Racz and Fitzgerald 2001), respectively. Approximately 90% of the total N existed in the form of NH<sub>4</sub>-N. Total N, P, K, S, Ca, and Mg contents and EC in the manure used in 2003 and 2004 were higher than that used in 2002, whereas total Na and pH were similar.

**Table 6. Levels of extractable NO<sub>3</sub>-N (µg/g) in soil samples collected at different times and depths.**

Year and factor*	Soil sampling time after injection and sampling depth											
	3 wk		5 wk		7 wk		9 wk		11 wk			
2002	0-300 mm		0-300 mm		0-300 mm		0-300 mm		0-300 mm			
<i>Tool type</i>												
Coulter	27.9a**		10.8a		6.6a		3.5a		1.6a			
Furrower	26.5a		13.1a		9.1a		2.9a		1.8a			
<i>Tool spacing</i>												
S300	32.3a		13.3a		9.3a		4.1a		2.2a			
S600	20.1b		8.4a		2.9b		1.9b		1.1b			
S900	29.4a		14.2a		11.1a		3.6a		1.8a			
2003	3 wk		6 wk		9 wk							
	0-300 mm		300-600 mm		0-300 mm		300-600 mm		0-300 mm		300-600 mm	
<i>Tool type</i>												
Coulter	25.6a		14.0a		10.9b		11.0a		16.6b		9.3a	
Furrower	28.4a		14.0a		17.5a		13.0a		20.9a		9.7a	
<i>Tool spacing</i>												
S300	29.8a		13.1b		14.1ab		11.8a		18.3ab		10.0a	
S600	25.8a		13.5b		11.8b		10.2a		16.7b		9.0a	
S900	25.5a		15.5a		16.6a		14.0a		21.2a		9.5a	
2004	3 wk		6 wk		19 wk (after harvest)							
	0-300 mm		300-600 mm		0-300 mm		300-600 mm		0-300 mm		300-600 mm	
<i>Tool type</i>												
Coulter	19.4a		19.0a		11.3b		12.6a		4.8a		2.1a	
Furrower	18.4a		17.5a		15.9a		14.2a		5.7a		2.7a	
<i>Tool spacing</i>												
S300	21.7a		21.0a		13.1a		14.7a		5.5a		3.21	
S600	20.7ab		18.7ab		14.4a		13.6ab		4.7a		1.8a	
S900	14.2b		15.0b		13.5a		11.9b		5.6a		2.2a	

\* S300, S600, and S900 refer to 300-, 600-, and 900-mm tool spacing treatments, respectively.

\*\* Mean values for each factor within a year and a column followed by the same letter are not significantly different ( $P > 0.1$ ).

### Extractable soil nutrient levels

**Soil nitrate nitrogen (NO<sub>3</sub>-N)** Soil NO<sub>3</sub>-N tended to be higher in plots where manure was applied using the furrower than in plots where manure was applied using the coulter (Table 6). However, there were no statistically significant differences observed at the 300-600 mm depth for all three years. At the 0-300 mm depth, use of the furrower rather than the coulter tended to result in higher levels of soil NO<sub>3</sub>-N; this difference was significant in two out of three samplings in 2003 and in one out of three samplings in 2004. This is in agreement with results observed from dairy and swine studies conducted by Schmitt et al. (1995). They reported that, in Minnesota, levels of soil NO<sub>3</sub>-N resulting from the use of winged tools were consistently higher than those from non-winged tools over the growing season. Similarly, Sawyer et al. (1990) reported increases in soil NO<sub>3</sub>-N when using winged tools rather than non-winged tools for applying liquid beef manure in Illinois. Schmitt et al. (1995)

suggested a twofold explanation for this observation. First, use of winged tools does not promote the levels of denitrification associated with using non-winged tools. Secondly, spatial distribution of manure might expedite mineralization of organic N as a result of shallower manure placement and increased contact between manure and soil.

Tool spacing also significantly affected the soil NO<sub>3</sub>-N. However, mixed results were observed among years perhaps due to differences in weather conditions. In 2002, levels of soil NO<sub>3</sub>-N for S300 and S900 were similar and higher by 77 and 74% (on average) than that for S600, respectively (Table 6). The reason why S600 resulted in lower NO<sub>3</sub>-N than both S300 and S900 is unknown. In 2003, soil NO<sub>3</sub>-N levels of S300 tended to be higher than those of S600 in both depth ranges, although they were not statistically significant (Table 6). Again, the significantly higher soil NO<sub>3</sub>-N levels for S900 were not explainable.

**Table 7. Levels of extractable NH<sub>4</sub>-N (µg/g) in soil samples collected at different times and depths.**

Year and factor*	Soil sampling time after injection and sampling depth			
	3 wk		6 wk	
	0-300 mm	300-600 mm	0-300 mm	300-600 mm
2002				
	3 wk	5 wk	7 wk	
	0-300 mm	0-300 mm	0-300 mm	
<i>Tool type</i>				
Coulter	0.47a**	1.12a	0.56a	
Furrower	0.48a	1.26a	0.54a	
<i>Tool spacing</i>				
S300	0.45a	1.44a	0.51a	
S600	0.52a	0.90b	0.57a	
S900	0.46a	1.23a	0.57a	
2004				
<i>Tool type</i>				
Coulter	17.3a	8.4a	10.8a	10.4a
Furrower	15.1a	8.5a	10.6a	10.6a
<i>Tool spacing</i>				
S300	13.4b	8.5a	11.7a	11.6a
S600	14.6b	8.2a	10.7a	9.9b
S900	20.6a	8.6a	9.7a	10.0b

\* S300, S600, and S900 refer to 300-, 600-, and 900-mm tool spacing treatments, respectively. Data were not collected in 2003.

\*\* Mean values for each factor within a year and a column followed by the same letter are not significantly different ( $P > 0.1$ ).

In 2004, the level of soil NO<sub>3</sub>-N in S300 was similar to that of S600, and the same was true between S600 and S900 at both depth ranges (Table 6). However, S300 had significantly higher soil NO<sub>3</sub>-N than S900 in one and two out of three sampling times at the 0-300 and 300-600 mm depths, respectively. This may be attributed to a possible greater denitrification loss in the S900 plots than in the S300 and S600 plots, favoured by a combination of large manure volume per manure band and the wetter soil condition in 2004.

With progress in the growing seasons, decreases in the overall soil NO<sub>3</sub>-N were observed due to uptake by plants. At the last sampling, levels of soil NO<sub>3</sub>-N were reduced by up to 95, 40, and 75% (maximum) of that at the first sampling following manure injection in 2002, 2003, and 2004, respectively, in the 0-300 mm soil depth.

**Soil ammonium nitrogen (NH<sub>4</sub>-N)** Analyses were not done for soil NH<sub>4</sub>-N on samples collected in 2003 due to a budget constraint. In comparison to 2002, more of the inorganic soil nitrogen was present in the form of NH<sub>4</sub>-N in 2004 (Table 7). In the dry year (2002), most of the manure NH<sub>4</sub>-N was possibly transformed to NO<sub>3</sub>-N but in the wet year (2004) that did not occur. Probably the cooler temperature and wetter condition in 2004 reduced nitrification.

The type of injection tool did not affect the level of soil NH<sub>4</sub>-N in 2002 and 2004 as indicated by the similar NH<sub>4</sub>-N

levels under both tools (Table 7). No particular trends were observed for spacing effects on the soil NH<sub>4</sub>-N levels. Three and seven weeks after injection, there were no significant differences in the levels of soil NH<sub>4</sub>-N between the tool spacing treatments in 2002. Five weeks after injection, soil NH<sub>4</sub>-N of S300 and S900 were higher by 60 and 37% than that of S600, respectively. In 2004, three weeks after injection, soil NH<sub>4</sub>-N increased with increasing tool spacing at the 0-300 mm depth; however it was not affected by the tool spacing at 300-600 mm depth. Six weeks after injection, soil NH<sub>4</sub>-N decreased with the tool spacing at the 300-600 mm depth.

Over the growing season in 2002, soil NH<sub>4</sub>-N during second sampling increased by more than 70% as compared to the first sampling, which may be due to net mineralization. At the third sampling, levels of soil NH<sub>4</sub>-N were back to their values at the time of first sampling. Similarly, in 2004 levels of soil NH<sub>4</sub>-N fluctuated over time. Again, this might be due to the combined effects of net mineralization and plant uptake.

**Soil phosphate (P<sub>2</sub>O<sub>5</sub>)** Significantly higher soil P<sub>2</sub>O<sub>5</sub> levels were observed for the furrower than for the coulter in 17 observations over the three years (Table 8). Tool spacing also significantly influenced the level of soil P<sub>2</sub>O<sub>5</sub>. In 2002, S900 resulted in a higher soil P<sub>2</sub>O<sub>5</sub> than S300 and S600 by 36 and 83%, respectively. This could be due to sampling errors. Eleven weeks after injection, S600 resulted in significantly lower level of P<sub>2</sub>O<sub>5</sub> than S300 and S900. This isolated observation was

also difficult to explain. In 2003, higher soil P<sub>2</sub>O<sub>5</sub> was observed in the S900 plots than in the S300 plots in both ranges of soil depth six weeks after manure injection. Over the growing season of 2004, soil P<sub>2</sub>O<sub>5</sub> of S300 was higher than those of S600 and S900. After harvest, however, levels of soil P<sub>2</sub>O<sub>5</sub> in the S300 and S900 plots were not significantly different.

**Potassium (K), sulphur (SO<sub>4</sub>-S), pH, and EC** Neither tool type nor the tool spacing affected levels of soil K, SO<sub>4</sub>-S, pH, and EC. Therefore, no detailed data are presented. Instead values of these variables averaged over the growing season of each year are summarized in Table 9.

### Crop performance

**Plant tillers, heads, main stem length, and above ground biomass** No significant differences were detected in plant number of tillers, heads, and main stem length between the furrower and the coulter in any of the three years. The furrower resulted in approximately 10% more plant biomass than the coulter in 2002 (Table 10); however, in 2003 and 2004, both tools yielded similar amounts of plant biomass.

In 2002, plant number of tillers, heads, main stem length, and above ground biomass followed a decreasing trend with increasing tool spacing (Table 10). This trend was significant for the biomass data among three tool spacing treatments. In 2003, there were no significant differences in any of the plant number of tillers, heads, main stem length, and biomass caused

**Table 8. Levels of extractable P<sub>2</sub>O<sub>5</sub> (µg/g) in soil samples collected at different times and depths.**

Year and factor*	Soil sampling time after injection and sampling depth					
	3 wk		6 wk		9 wk	
2002	0-300 mm		0-300 mm		0-300 mm	
<i>Tool type</i>						
Coulter	101.1a**		84.5a		139.3a	
Furrower	106.1a		91.0a		126.2a	
<i>Tool spacing</i>						
S300	101.1a		84.5a		128.7b	
S600	101.6a		87.0a		95.5b	
S900	107.6a		91.5a		174.5a	
2003	3 wk		6 wk		9 wk	
	0-300 mm		300-600 mm		0-300 mm	
	300-600 mm		0-300 mm		300-600 mm	
<i>Tool type</i>						
Coulter	12.9a		4.7b		10.5a	
Furrower	13.0a		5.7a		11.6a	
<i>Tool spacing</i>						
S300	13.1a		5.1a		10.0b	
S600	13.1a		5.2a		11.4ab	
S900	12.7a		5.1a		11.8a	
2004	3 wk		6 wk		19 wk (after harvest)	
	0-300 mm		300-600 mm		0-300 mm	
	300-600 mm		0-300 mm		300-600 mm	
<i>Tool type</i>						
Coulter	77.7a		53.6a		59.4a	
Furrower	73.7a		54.7a		61.1a	
<i>Tool spacing</i>						
S300	76.0a		64.3a		69.3a	
S600	72.3a		51.5b		55.9b	
S900	78.9a		46.7b		55.6b	

\* S300, S600, and S900 refer to 300-, 600-, and 900-mm tool spacing treatments, respectively.

\*\* Mean values for each factor within a year and a column followed by the same letter are not significantly different (P > 0.1).

**Table 9. Average values of extractable soil K (µg/g), SO<sub>4</sub>-S (µg/g), pH, and EC (dS/m) in different years at different sampling depths.**

Factor*	K			SO <sub>4</sub> -S	pH			EC		
	2002	2003		2002	2002	2004		2002	2004	
	0-300 mm	0-300 mm	300-600 mm	0-300 mm	0-300 mm	0-300 mm	300-600 mm	0-300 mm	0-300 mm	300-600 mm
<i>Tool type</i>										
Coulter	482	269	224	8	7.68	7.54	8.06	0.63	0.59	0.70
Furrower	479	279	229	10	7.70	7.47	7.93	0.61	0.56	0.81
<i>Tool spacing</i>										
S300	484	267	221	9	7.71	7.53	7.97	0.66	0.60	0.74
S600	463	275	231	9	7.71	7.50	8.04	0.58	0.55	0.65
S900	495	279	228	10	7.64	7.48	7.97	0.62	0.57	0.87

\* S300, S600, and S900 refer to 300-, 600-, and 900-mm tool spacing treatments, respectively.

**Table 10. Plant number of tillers, heads, main stem length, and above ground biomass.**

Year and factor*	No. of tillers	No. of heads	Main stem length (mm)	Biomass (kg/ha)
2002				
<i>Tool type</i>				
Coulter	2.8a**	2.5a	582a	6702b
Furrower	3.1a	2.8a	572a	7339a
<i>Tool spacing</i>				
S300	3.1a	2.7a	587a	7545a
S600	3.0a	2.7a	573a	6853b
S900	2.8a	2.5a	571a	6663b
2003				
<i>Tool type</i>				
Coulter	2.2a	1.8a	510a	4445a
Furrower	2.3a	1.9a	512a	4390a
<i>Tool spacing</i>				
S300	2.3a	1.9a	500a	4322a
S600	2.3a	1.9a	516a	4446a
S900	2.1a	1.7a	517a	4484a
2004				
<i>Tool type</i>				
Coulter	7.2a	6.2a	959a	8471a
Furrower	7.2a	6.6a	972a	8355a
<i>Tool spacing</i>				
S300	7.9a	7.2a	978a	8473a
S600	6.6b	5.9b	949b	8306a
S900	7.1ab	6.1b	969a	8459a

\* S300, S600, and S900 refer to 300-, 600-, and 900-mm tool spacing treatments, respectively.

\*\* Mean values for each factor within a year and a column followed by the same letter are not significantly different ( $P > 0.1$ ).

by the tool spacing. In 2004, S300 resulted in higher plant number of tillers and heads and longer main stem than S600 and S900. There were no differences in above ground biomass between the spacing treatments in that year.

**Grain and straw yields** It is obvious that grain yields were dictated by the weather conditions. Both grain and straw yields were the lowest in the driest growing season of 2003 and they were highest in the wettest growing season (2004) (Table 11). The low yield in 2003 may be explained by the fact that the plant nutrient uptake was undermined by dry weather conditions (Table 11).

No significant differences were detected in grain yield between the furrower and the coulter (Table 11). Tool spacing did not significantly affect grain and straw yields in any of the growing seasons. Differences in grain yield due to injection tool type and spacing may have been masked due to the effect of late seeding on grain yield. Sawyer et al. (1991) reported inconsistent results in grain yield among tool types. Schmitt et al. (1995) observed higher grain yield when using winged tools than non-winged tools.

Warner and Godwin (1988) examined grass response to injected sewage sludge at various injector tool spacings and found that a 650-mm tool spacing resulted in higher grass yield than 500 and 850-mm tool spacings. Eghball and Sander (1989) studied band spacing effects of dual-placed N and P fertilizers on corn. Their observation was similar to the results of this study in that band spacing did not affect corn yield unless either N or P deficiency dominated the total input of the dual placed band. Results of this study agree with findings of Maxwell et al. (1984) who reported that 250 and 380-mm spacings resulted in more uniform plant growth and dry matter production early in the growing season, but the effects on yield were not significant, when compared to 500-mm spacing.

**Total N and P in plant biomass, grain, and straw** In 2002, as compared with the coulter, the furrower resulted in higher levels of total N in the plant biomass, and total N and P in grain and straw (Table 12). In 2003 and 2004, the amounts of total N and P in plant biomass, grain, and straw were similar when manure was applied using either the coulter or the furrower. One exception was that, in 2004, 10% higher total N and 21% lower P in straw were measured when using the furrower rather than the coulter.

Tool spacing did not significantly affect total N and total P in plant materials in 2002 (Table 12). In 2003 and 2004, S300 had higher plant nutrient values than S600 and S900. Statistically significant differences were observed for total N and total P in grain in 2003 and for total N in biomass in 2004.

## CONCLUSIONS

Compared to the coulter-type, the furrower-type injection tool offered a slight advantage in terms of increased soil nitrate, plant biomass production, total N concentration in biomass, grain and straw, and total P in grain and straw. Among the tool spacings of 300, 600, and 900 mm, the 300-mm spacing resulted in higher levels of soil nitrate. Although the narrowest injection tool spacing did not offer any advantage over the other tool spacings in terms of yield response, better plant development and higher plant biomass production were observed for the 300-mm tool spacing, which is important with regard to nutrient cycling in agricultural systems. Considering the resultant higher levels of soil nitrate and better crop performance, the furrower-type tool spaced 300-mm apart was the best choice for liquid manure injection in spring barley production. It would be of particular interest to perform similar experiments on different soil types, and under varying climatic conditions, to confirm the observed trends since this experiment was carried out on heavy clay-loam lacustrine soils under less than optimum growing conditions.

## ACKNOWLEDGEMENTS

We gratefully acknowledge the financial support from the Natural Sciences and Engineering Research Council of Canada (NSERC) and Manitoba Agriculture, Food, and Rural Initiatives (Covering New Ground program). We also appreciate assistance from Dr. Sylvio Tessier, Dr. Shafiqur Rahman, Razif Mahadi, Faustinus Monero, Van Doan, Erron Leafloor, Dee Genaille, Gerry Woods, and Randy Westwood in the fieldwork.



**Table 11. Grain and straw yields for different treatments in three years.**

Factor*	Grain yield (kg/ha)			Straw yield (kg/ha)		
	2002	2003	2004	2002	2003	2004
<i>Tool type</i>						
Coulter	2959a**	1229a	3744a	1776a	1194a	4702b
Furrower	2850a	1286a	3612a	1712a	1207a	5089a
<i>Tool spacing</i>						
S300	2896a	1276a	3522a	1713a	1271a	4962a
S600	2893a	1306a	3837a	1773a	1181a	4977a
S900	2924a	1190a	3674a	1745a	1150a	4748a

\* S300, S600, and S900 refer to 300-, 600-, and 900-mm tool spacing treatments, respectively.

\*\* Mean values for each factor within a year and a column followed by the same letter are not significantly different ( $P > 0.1$ ).

**Table 12. Total N (TN) and P (TP) in plant biomass, grain, and straw.**

Year and factor*	Biomass		Grain		Straw	
	TN (%)	TP (%)	TN (%)	TP (%)	TN (%)	TP (%)
2002						
<i>Tool type</i>						
Coulter	1.32b**	0.18a	1.91b	0.40b	0.93b	0.15b
Furrower	1.50a	0.17a	1.97a	0.41a	1.08a	0.17a
<i>Tool spacing</i>						
S300	1.40a	0.17a	1.96a	0.40a	1.03a	0.16a
S600	1.47a	0.18a	1.93a	0.40a	1.00a	0.15a
S900	1.37a	0.18a	1.93a	0.41a	0.97a	0.16a
2003						
<i>Tool type</i>						
Coulter	1.84a	0.11a	2.30a	0.32a	1.41a	0.05a
Furrower	1.90a	0.12a	2.25a	0.32a	1.40a	0.05a
<i>Tool spacing</i>						
S300	1.95a	0.12a	2.35a	0.34a	1.46a	0.06a
S600	1.83a	0.12a	2.22b	0.31b	1.41a	0.05a
S900	1.82a	0.11a	2.25b	0.32ab	1.36a	0.05a
2004						
<i>Tool type</i>						
Coulter	2.5a	0.14a	2.0a	0.29a	1.37b	0.14a
Furrower	2.5a	0.14a	2.0a	0.31a	1.50a	0.11b
<i>Tool spacing</i>						
S300	2.6a	0.14a	2.1a	0.31a	1.47a	0.13a
S600	2.5ab	0.15a	2.0a	0.30a	1.40a	0.13a
S900	2.4b	0.14a	2.0a	0.30a	1.44a	0.12a

\* S300, S600, and S900 refer to 300-, 600-, and 900-mm tool spacing treatments, respectively.

\*\* Mean values for each factor within a year and a column followed by the same letter are not significantly different ( $P > 0.1$ ).

**REFERENCES**

- Alberta Research Council. 1996. Methods manual for chemical analysis of water and wastes. Method 19103 565, Potassium, dissolved, automated photometry method. Edmonton, AB: Alberta Research Council.
- AOAC. 1990. *Official Methods of Analysis of AOAC*, 15<sup>th</sup> edition. Arlington, VA: Association of Official Analytical Chemists International.
- ASAE. 2004a. ASAE S358.2 Moisture measurement-Forages. In *ASAE Standards 2004*, 584. St. Joseph, MI: ASABE.
- ASAE. 2004b. ASAE S414.1 FEB04. Terminology and definitions for agricultural tillage implements. In *ASAE Standards 2004*, 274-285. St. Joseph, MI: ASABE.
- Ashworth, J. and Mrazek K. 1995. "Modified Kelowna" test for available phosphorus and potassium in soil. *Communications in Soil Science and Plant Analysis* 26(5&6): 731-739.
- Chen, Y. and R. Samson. 2002. Integration of liquid manure into conservation tillage corn systems. *Transactions of the ASAE* 45(3): 629-638.
- Chen, Y. and S. Tessier. 2001. Criterion for design and selection of injection tools to minimize liquid manure on the soil surface. *Transactions of the ASAE* 44(6): 1421-1428.
- Clesceri, L.S., A.E. Greenbert, A.D. Eaton and M.A.H. Franson. 1998. *Standard Methods for the Examination of Water and Wastewater*, 20<sup>th</sup> edition. Washington, DC. American Public Health Association.
- Comfort, S.D., K.A. Kelling, D.R. Keeney and J.C. Converse. 1988. The fate of nitrogen from injected liquid manure in a silt loam soil. *Journal of Environmental Quality*. 17: 317-322.
- Eghball, B. and D.H. Sander. 1989. Band spacing effects of dual-placed nitrogen and phosphorus fertilizers on corn. *Agronomy Journal* 81(2): 178-184.

- Fitzmaurice, J., E.S. Jacques, A. Waddell and R. Eilers. 1999. Soils of the Brandon Research Centre – updated from Hopkins and St. Jacques, 1985. Special Report Series 99-2. Brandon, MB: Land Resource Unit, Brandon Research Centre, Agriculture and Agri-Food Canada.
- Greenberg, A.E., L.S. Clesceri and A.D. Eaton. 1992. *Standard Methods for Examination of Water and Wastewater*, 18<sup>th</sup> edition. Washington, DC: American Public Health Association.
- Grevers, M. and J. Schoenau. 2001. Swine manure - a dilute fertilizer with extra's. In *Proceedings of the 2001 Soils & Crops Workshop*, 452-456. Saskatoon, SK: University of Saskatchewan.
- Hamm, J.W., F.G. Radford and E.H. Halstead. 1970. The simultaneous determination of nitrogen, phosphorus and potassium in sodium bicarbonate extracts of soils. In *Advances in Automatic Analysis, Vol. 2: Industrial Analysis Technicon International Congress*, 65-69. Orlando, FL: Futura Publishing.
- Hendershot, W.H., H. Lalonde and M. Duquette. 1993. Soil reaction and exchangeable acidity. In *Soil Sampling and Methods of Soil Analysis*, ed. M.R. Carter, 141-144. Boca Raton, FL: CRC Press.
- Hoff, J.D., D.W. Nelson and A.L. Sutton. 1981. Ammonia volatilization from liquid swine manure applied to cropland. *Journal of Environmental Quality* 10(1): 90-95.
- Maxwell, T.M., D.E. Kissel, M.G. Waggoner, D.A. Whitney, M.L. Cabrera and H.C. Moser. 1984. Optimum spacing of preplant bands of N and P fertilizer for winter wheat. *Agronomy Journal* 76(2): 243-247.
- Maynard, D.G. and Y.P. Kalra. 1993. Nitrate and exchangeable ammonium nitrogen. In *Soil Sampling and Methods of Soil Analysis*, ed. M.R. Carter, 25-38. Boca Raton, FL: CRC Press.
- McCormick, R.A., D.W. Nelson, A.L. Sutton and D.M. Huber. 1983. Effect of nitrapyrin on nitrogen transformation in soil treated with liquid swine manure. *Agronomy Journal* 75(6): 947-950.
- McKeague, J.A. 1976. Soluble salts. Manual on soil sampling and methods of analysis. Subcommittee of Canada Soil Survey Committee. Ottawa, ON: Agriculture and Agri-Food Canada.
- Mooleki, S.P., J.J. Schoenau, G. Hultgreen, G. Wen and J.L. Charles. 2002. Effect of rate, frequency and method of liquid swine manure application on soil nitrogen availability, crop performance and N use efficiency in east-central Saskatchewan. *Canadian Journal of Soil Science* 82(4): 457-467.
- Olsen, S.R. and L.E. Sommers. 1982. Phosphorus. In *Methods of soil analysis Part 2*, ed. A.L. Page, 403-430. Madison, WI: American Society of Agronomy and Soil Science Society of America.
- Qian, P., J.J. Schoenau and R.E. Karamanos. 1994. Simultaneous extraction of available phosphorus and potassium with a new soil test: A modification of the Kelowna extraction. *Communications in Soil Science and Plant Analysis* 25(5&6): 627-635.
- Racz, G.J. and M.M. Fitzgerald. 2001. Nutrient and heavy metal contents of hog manure – effects on soil quality and productivity. In *Livestock Options for the Future: A National Conference on Livestock & the Environment*, <http://www.gov.mb.ca/agriculture/livestock/livestockopt/papers/racz.pdf> (2006/11/03).
- Rahman, S., Y. Chen, K. Buckley and W. Akinremi. 2004. Slurry distribution in soil as influenced by slurry application micro-rate and injection tool type. *Biosystems Engineering* 89(4): 495-504.
- SAS Institute Inc. 2001. SAS STAT7 User's Guide, Version 8.2. Cary, NC: SAS Institute, Inc.
- Sawyer, J.E., M.A. Schmitt and R.G. Hoefl. 1990. Inorganic nitrogen distribution and soil chemical transformations associated with injected liquid beef manure. *Agronomy Journal* 82(5): 963-969.
- Sawyer, J.E., M.A. Schmitt, R.G. Hoefl, J.C. Siemens and D.H. Vanderholm. 1991. Corn production associated with liquid beef manure application methods. *Journal of Production Agriculture* 4(3): 335-344.
- Schmitt, M.A., S.D. Evans and G.W. Randall. 1995. Effect of liquid manure application methods on soil nitrogen and corn grain yields. *Journal of Production Agriculture* 8(2): 186-189.
- Sutton, A.L. 1994. Proper animal manure utilization. *Journal of Soil and Water Conservation* 49(2): 65-70.
- Thomas, R.L., R.W. Sheard and J.R. Moyer. 1967. Comparison of conventional and automated procedures for nitrogen, phosphorus, and potassium analysis of plant material using a single digestion. *Agronomy Journal* 59: 240-243.
- Warner, N.L. and R.J. Godwin. 1988. An experimental investigation into factors influencing the soil injection of sewage sludge. *Journal of Agricultural Engineering Research* 39(4): 287-300.